

A Meta-Analysis Shows That Docosahexaenoic Acid from Algal Oil Reduces Serum Triglycerides and Increases HDL-Cholesterol and LDL-Cholesterol in Persons without Coronary Heart Disease^{1–3}

Adam M. Bernstein,^{4,5*} Eric L. Ding,^{5,6} Walter C. Willett,^{5–7} and Eric B. Rimm^{5–7}

⁴Wellness Institute of the Cleveland Clinic, Lyndhurst, OH; ⁵Department of Nutrition, and ⁶Department of Epidemiology, Harvard School of Public Health, Boston, MA; and ⁷Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

Abstract

Certain algae contain the (n-3) fatty acid DHA, yet the relation between algal oil supplementation and cardiovascular disease risk factors has not been systematically examined. Our objective was to examine the relation between algal oil supplementation and cardiovascular disease risk factors. We conducted a systematic review of randomized controlled trials published between 1996 and 2011 examining the relation between algal oil supplementation and cardiovascular disease risk factors and performed a meta-analysis of the association between algal oil DHA supplementation and changes in the concentrations of TG, LDL-cholesterol (LDL-C), and HDL-cholesterol (HDL-C). We identified 11 randomized controlled trials with 485 healthy participants that evaluated the relation between algal oil DHA supplementation and TG, LDL-C, and HDL-C. The median dose of algal DHA was 1.68 g/d. The pooled estimate for the change in TG concentration was -0.20 mmol/L (95% CI: -0.27 to -0.14), 0.23 mmol/L (95% CI: 0.16 – 0.30) for LDL-C, and 0.07 mmol/L (95% CI: 0.05 – 0.10) for HDL-C. **DHA supplementation from algal oil, a marine source of (n-3) fatty acids not extracted from fish, may reduce serum TG and increase HDL-C and LDL-C in persons without coronary heart disease.** J. Nutr. doi: 10.3945/jn.111.148973.

Introduction

Consumption of fatty fish high in EPA and DHA is recommended for healthy persons and those with high blood pressure, elevated TG, and CHD⁸ (1–4). There may also be benefit for persons with heart failure (5). Consumption of wild marine fish is severely limited, however, because most ocean fish populations are exploited and unable to yield sustainable catches at current levels (6–8). Aquaculture, or fish farming, provides an alternative or perhaps additional source of fish for consumption, yet the practice has been associated with damage to and pollution of sensitive ecologic areas (8–10).

In the wild, fish accumulate EPA and DHA by consumption of marine algae (11,12). Recently, algae have been cultivated to preferentially produce DHA and their oil has been extracted to make supplements (13,14). Yet although narrative reviews have

been conducted on the relation between algal oil supplements and CHD risk factors, such as serum lipids (15,16), there have not yet been systematic examinations. Given the clinical, public health, and ecologic importance of understanding algal oil supplementation, we performed a systematic review and meta-analysis of published, randomized, controlled clinical trials that evaluated and quantified the association between algal oil and cardiovascular disease risk factors.

Materials and Methods

We observed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines for this systematic review and meta-analysis (17).

Search strategy. We searched for all randomized controlled trials that examined the association between algae-derived EPA or algae-derived DHA and serum lipoproteins or cardiovascular disease. We searched the following online databases through May 12, 2011: MEDLINE, EMBASE, CABI (CAB abstracts), Cochrane Central Registry of Controlled Trials, www.clinicalstudyresults.org (a Web-based database of results from trials of U.S.-marketed pharmaceuticals), and www.clinicaltrials.gov (a registry of federally and privately supported clinical trials in the U.S. and around the world) (The search strategy is available in the online material). There were no language restrictions in our search.

¹ Supported by the Harvard Human Nutrition Program (A.M.B.).

² Author disclosures: A. M. Bernstein, E. L. Ding, W. C. Willett, and E. B. Rimm, no conflicts of interest.

³ Supplemental Tables 1–5 and Supplemental Methods are available from the "Online Supporting Material" link in the online posting of the article and from the same link in the online table of contents at <http://jn.nutrition.org>.

⁸ Abbreviations used: CHD, coronary heart disease; HDL-C, HDL-cholesterol; LDL-C, LDL-cholesterol.

* To whom correspondence should be addressed. E-mail: bernsta2@ccf.org.

We excluded studies of animals and children and of persons with diseases that may affect absorption or metabolism of algal oil (e.g., patients with cystic fibrosis).

Study selection. Our search spanned all published studies prior to May, 2011. In our first round of searching in May, 2010, we initially located 756 articles: 185 articles from MEDLINE, 201 from EMBASE, 317 from CABI, and 53 from Cochrane Central Registry of Controlled Trials. We did not identify any relevant completed trials from www.clinicaltrials.gov. An additional MEDLINE search for articles written by select authors and a review of their bibliographies identified an additional 9 articles. A total of 736 articles were excluded after reviewing the title or abstract. Full texts of the remaining 29 were selected for secondary review. In May, 2011, we conducted an updated search for newly published articles and identified one additional article. We did not identify any studies that included cardiovascular disease outcomes. We identified studies that evaluated algal oil supplementation on thrombosis (18,19), inflammatory markers (19–22), blood pressure (19,21–23), plasma lipids (FFA) (18–20,22,24–31), and erythrocyte membrane lipids (19,20,24,26,27,30–32), but given the potential for unstable estimates from a limited amount of data due to small number of participants in these studies, we did not include these endpoints in our meta-analysis and instead focused on the relation between algal oil supplementation and serum lipids. We included 11 unique studies in the final analysis (18,19,22,23,26,28,29,33–36).

Data extraction and quality assessment. For each of the final 11 studies, we reviewed and extracted the following data using a piloted form: first author and study year, number of study participants, mean age of study participants, sex distribution of participants, study design (randomization, blinding, presence of placebo, parallel or crossover design), dose of EPA or DHA in the algal oil supplement as well as other fatty acids, content of placebo, duration of intervention, whether the study was funded by industry, and whether the algal oil was gifted to the investigators by industry and, if so, whether there was independent assessment of the supplement content. We examined whether each study used fasting blood samples for measuring lipid concentration, whether the LDL-C concentration was derived with the Friedewald equation (37), and if LDL particle size was estimated. The principal study outcomes extracted were the mean and SEM of the treatment effect of algal oil supplementation on TG, HDL-C, and LDL-C. For studies that did not directly report the treatment effect, we calculated it from the difference in the within-group estimates of change in lipid concentration (18,22,23,29,33,34,36) or from the difference in the final mean concentration of lipid concentrations (28). For studies that did not directly report the variance, we calculated it from data provided (Supplemental Methods). One study reported two intervention groups with one control group (29); for our analysis, we collapsed the two intervention groups into one group and calculated an average intervention dose and average treatment effect to avoid statistical duplication in the meta-analysis. All algal oil supplements used in the analyzed studies contained either no EPA (18,22,23,29,33,35,36) or a negligible amount of EPA (19,26,28,34) (e.g., 1/20 the amount of DHA/d or <0.005 g/d). To convert mg/dL of TG to mmol/L, we divided by 89; to convert mg/dL of HDL-C or LDL-C to mmol/L of HDL-C or LDL-C, we divided by 39 (38).

The quality of each study, defined as the confidence that its design, conduct, analysis, and presentation limited bias in its results, was assessed by a validated quality scoring system (39). The scoring system awarded points to each study based on: 1) the presence of randomization; 2) appropriate method of randomization (e.g., using random numbers); 3) presence of double-blinding; 4) appropriate method of double-blinding (e.g., using supplements with similar appearance); and 5) description of withdrawals and dropouts. The lowest possible quality score was 0 and the maximum was 5.

Statistical analysis. To derive a summary (pooled) estimate of the association between algal oil DHA supplementation (g/d) and serum TG, LDL-C, and HDL-C concentrations (mmol/L), we performed a fixed-effects meta-analysis of the treatment effect, which was equal to the weighted mean difference in serum lipid concentration between inter-

vention and control groups (*metan* procedure, STATA 11, StataCorp). The fixed-effects model assumes that the true effect of treatment is the same in each study and that differences between study results are due to chance (40). Study weights were determined by the inverse variance method so that the weight given to a particular study was the reciprocal of the squared SEE estimate. We derived an estimate of the pooled percent change in TG, LDL-C, and HDL-C due to algal oil supplementation by dividing the weighted pooled absolute change by the mean baseline lipid concentration (mean baseline concentration equals the sum of mean baseline concentration in each study multiplied by its fixed-effects weight). Magnitude of between-study heterogeneity was examined by determination of the Q statistic and I-squared statistic (41). Forest plots were used to visually graph the association between algal oil DHA supplementation and changes in absolute concentrations of serum TG, HDL-C, and LDL-C.

In sensitivity analyses to evaluate for potential dose-dependent effects, we fit three separate fixed-effects meta-regression models in which the outcomes were continuous variables for the treatment effect on TG, LDL-C, or HDL-C and the independent variable was mean algal DHA dose (g/d) (*vuuls* procedure, STATA 11, StataCorp). In these meta-regression dose-effect models, the coefficient for the algal oil variable is an estimate of the dose-response relation between algal oil and change in serum lipid concentration.

Potential for publication bias was assessed using the Begg rank correlation method and Egger weighted regression test (40) for serum TG, LDL-C, and HDL-C and by inspecting three funnel plots of treatment effect compared to SE for TG, LDL-C, and HDL-C. Trim-and-fill techniques were used in sensitivity analyses to correct for potential publication bias. $P < 0.05$ was considered significant.

Results

We identified 11 randomized controlled trials, including a total of 485 participants, dating from 1996 to 2011 (18,19,22,23,26,28,29,33–36) (Supplemental Table 1). The mean age of study participants ranged from 24 to 59 y. Most studies were 6 wk in duration and either did not independently assess algal oil concentration or did not report such an assessment (6 of 11 studies). Every study either received the algal oil as a gift from industry or the study itself was funded by industry. All studies used fasting blood samples for lipid measurements and the majority used the Friedewald equation to estimate LDL-C. All but one of the studies reported that the algal oil contained a mixture of SFA and PUFA in addition to DHA, although the overall mean DHA fatty acid content was 41% by weight (Supplemental Table 2). Studies were of good quality (9 of 11 scored 4 of 5).

The median dose of algal DHA in the analysis of serum lipids was 1.68 g/d. The pooled estimate for the change in TG concentration was -0.20 mmol/L (95% CI: -0.27 to -0.14) (Fig. 1 and Supplemental Table 3), for LDL-C it was 0.23 mmol/L (95% CI: 0.16 – 0.30) (Fig. 2 and Supplemental Table 4), and for HDL-C it was 0.07 mmol/L (95% CI: 0.05 – 0.10) (Fig. 3 and Supplemental Table 5). A 0.20 -mmol/L absolute reduction in TG corresponded to a 15% (95% CI: 10–20%) decrease, a 0.23 -mmol/L increase in LDL-C corresponded to an 8% (95% CI: 6–10%) increase, and a 0.07 -mmol/L increase in HDL-C corresponded to a 5% (95% CI: 4–7%) increase.

The greatest decrease in TG and greatest increase in LDL-C appeared in the two studies that included participants with elevated serum lipid concentrations at baseline (23,29). The two studies that showed a decrease in LDL-C included vegetarian participants (18,35), although the CI were wide, and a third study with vegetarian participants (34) observed a significant rise in LDL-C, similar to other studies. One study that reported using direct methods for estimating LDL-C, rather than the Friedewald equation, showed a decrease in LDL-C (35), whereas




FIGURE 1 Change in serum TG (mmol/L) associated with algal DHA supplementation (g/d). Weighted mean difference in absolute change estimated; median dose of algal oil = 1.68 g/d; I-squared (variation in effect size attributable to heterogeneity) = 0.6%; Q statistic *P* value = 0.43; *n* = 11.

another that used the direct method showed no effect (33). Limiting the study to the seven studies that reported using the Friedewald equation (18,19,22,23,28,29,34), the effect estimate of change in LDL-C was 0.29 (95% CI: 0.20–0.38) mmol/L LDL-C. One study reported using “standard” methods for estimating LDL-C (36) and one did not report its method (26). The one study that showed a higher increase in HDL-C compared to other studies had young study participants, was of longer duration than most studies, and was single-blinded (33).

There was little evidence of between-study heterogeneity by Q statistic and I-squared statistic; of the total variation in each weighted mean difference model, the amount of between-study variation was 0.6% when evaluating the association between algal DHA and TG (Q statistic *P* value = 0.43), 41% for algal oil and LDL-C (Q statistic *P* value = 0.08), and 31% for algal DHA and HDL-C (Q statistic *P* value = 0.15).




FIGURE 2 Change in serum LDL-C (mmol/L) associated with algal DHA supplementation (g/d). Weighted mean difference in absolute change estimated; median dose of algal oil = 1.68 g/d; I-squared (variation in effect size attributable to heterogeneity) = 40.8%; Q statistic *P* value = 0.08; *n* = 11. LDL-C, LDL-cholesterol.

In the meta-regression, we did not observe a significant dose response between algal DHA and TG, LDL-C, or HDL-C concentration: a 1-g increase per day in the dose of algal DHA was associated with a 0.06-mmol/L decrease in TG (95% CI: -0.19, 0.07), a 0.04-mmol/L reduction in LDL-C (95% CI: -0.28, 0.19), and a 0.02-mmol/L reduction in HDL-C (95% CI: -0.07, 0.04). All studies but one had doses of algal oil DHA <3 g/d. The magnitude of TG reduction was relatively constant between 0.7 and 2.8 g/d of algal DHA (Fig. 4). In the one study that gave 3 g/d of algal oil DHA, there was a further reduction in TG concentration (23).

The three funnel plots were symmetrical, suggesting that effect estimates were not associated with SE or study size. Of the three Begg rank correlation tests and three Egger tests, five were not significant and only the Egger test with serum TG was significant (*P* value 0.01), suggesting an overall small potential for publication bias or small-study effects. Trim-and-fill correction analysis with serum TG showed the main results remained significant and robust.

Discussion

To our knowledge, this is the first systematic review and meta-analysis of algae-derived DHA supplementation and serum lipids. We observed a significant decrease in TG and significant increases in both HDL-C and LDL-C. The 0.20-mmol/L (95% CI: -0.27 to -0.14) reduction in TG we observed with the 1.68-g/d dose of algal DHA is on the same order of magnitude observed with fish oil supplementation in a recent meta-analysis: 3.25 g of fish oil (including 1.90 g EPA and 1.35 g DHA) was associated with a 0.34-mmol/L (95% CI: -0.41 to -0.27) reduction in TG (42). The increase in HDL-C was also similar to that with fish oil (0.01 mmol/L; 95% CI: 0.00–0.02); the increase in LDL-C was higher (0.06 mmol/L; 95% CI: 0.03–0.09) (42).

Current AHA dietary recommendations for persons without CHD are to eat 2 servings/wk of fish (preferably oily) to reach ~0.25 g/d of the (n-3) fatty acids EPA and DHA and to reduce the risk of sudden death and death from CHD (3,43,44). The NIH has suggested 0.65 g/d of EPA and DHA and a minimum of 0.3 g/d of DHA during pregnancy and lactation (45), whereas the Institute of Medicine makes no specific recommendations for EPA and DHA intake (46). For persons with CHD, the AHA recommends 1 g/d of EPA and DHA, preferably from fish, but supplements can be considered in consultation with a physician. For patients with high TG, the recommendation is 2–4 g/d of EPA and DHA provided as capsules and under the supervision of a physician (3).

Recent data from the NHANES in 2005–2006 shows that U.S. citizens consume on average 0.05 g/d of EPA and 0.10 g/d of DHA (47), far less than that recommended either for healthy persons or individuals with high TG or CHD. Yet consumption of more fish, or extraction of their oil for supplements, may not be possible given the limitations of marine fishing. Stocks of Pacific salmon, tuna, herring, mackerel, and anchovies are already fully exploited (the ocean fishery is operating at or near optimal yield with no room for further expansion) or over-exploited (there is no room for further expansion and there is a high risk for depletion/collapse of the fish stock) (6,48,49) and Atlantic salmon stocks are also fully exploited to depleted (48). Although aquaculture provides an alternative to marine fishing, there are concerns with its ecologic impact (8–10).

Fish naturally accumulate EPA and DHA by consumption of marine algae (11,12), which are a diverse group of uni- and multi-cellular photosynthetic organisms. In contrast to algae, such as seaweed, microscopic algae have been commercially




FIGURE 3 Change in serum HDL-C (mmol/L) associated with algal DHA supplementation (g/d). Weighted mean difference in absolute change estimated; median dose of algal oil = 1.68 g/d; I-squared (variation in effect size attributable to heterogeneity) = 31.4%; Q statistic P value = 0.15; n = 11. HDL-C, HDL-cholesterol.

cultivated in open ponds and closed photo-bioreactors where they preferentially produce one type of PUFA (14). Algae currently being bred in this algaculture to create DHA-rich algal oil include *Cryptocodinium cohnii*, *Schizochytrium* species, and *Ulkenia* species. Food products already fortified with algal oil include infant formula, olive and canola oils, and soy milk (50).

Purified ethyl ester DHA supplementation has been shown to lower TG and mildly elevate LDL-C and HDL-C (51,52). The 8% increase in LDL-C and 5% increase in HDL-C we observed with algal oil is similar to that observed with 4 g/d of purified DHA, the latter which was also associated with a transition to larger, possibly less-atherogenic LDL particles (52). DHA is thought to decrease TG levels by reducing hepatic VLDL synthesis (53). Reduced synthesis in turn leads to reduced secretion and smaller VLDL particles, which are more readily converted to LDL than are large VLDL particles and which also compete with LDL for uptake by LDL receptors (52). Both of these actions lead to higher serum LDL-C. The mechanism by

which DHA increases HDL-C is not known but may be through decreased lipid transfer protein activity (54).

All but one of the placebo arms in the trials we examined contained corn, soy, or olive oil, each of which is rich in palmitic, oleic, and linoleic acids. Soy oil also has a small amount of α -linolenic acid. Most of the algal oil supplements contained palmitic and oleic acid, though in different amounts than the placebos. The DHA content of the supplements could explain our findings; however, it is also likely that the saturated fat content of the algal oil contributed to study participant lipid concentrations different than that those expected from purified DHA. Of the six studies that measured LDL particle size in relation to algal oil supplementation, five reported an increase in particle size and one reported no change; because particle size, in addition to lipoprotein cholesterol content, may help predict atherogenic risk, the net effect of algal DHA supplementation on serum lipoproteins and lipids may be beneficial despite the increase in LDL-C (55). We also note that two earlier meta-analyses of fish oil supplementation reported increase in LDL similar to those we observed in this study with algal oil (56,57). The effect of algal oil supplementation on overall cardiovascular risk must also consider its effect on other cardiovascular disease risk factors, such as blood pressure, platelet aggregation, and inflammation.

We did not identify any reports on algal oil contamination but, according to the U.S. Dietary Supplement and Health Education Act of 1994, the FDA cannot vet supplement content before a supplement is sold. Some fish oil supplements have been found to be contaminated with persistent organic pollutants (58,59). High-dose algal oil supplementation, as with fish oil supplementation, should be done under a physician's supervision to monitor for side effects, adverse reactions, and interactions with other medications. To approximate 200 mg/d of DHA, algal oil supplementation at present may cost three times as much as fish oil supplementation (60,61).

The strengths of our analysis are that we made extensive efforts to locate all published and unpublished (i.e., abstract only) work without limitation to language, country, date, or funding. Study designs were of good quality and with an average duration of 6 wk, there was adequate time to see changes in measured outcomes. The chief limitations are that the studies were few and small, all were funded or gifted by industry, and most did not independently verify the algal oil supplements. We were also only able to assess the relation between algal oil supplementation and serum lipids. Because one of the recognized benefits of fish oil supplementation is antiarrhythmic with a reduction in sudden death, our ability to compare algal oil supplementation to fish oil supplementation is limited. Moreover, dose-response results from the meta-regression should be interpreted cautiously, because the highest dose came from a study population with high baseline TG. Thus, residual between-study confounding is a potential limitation of our dose-response analysis. We included studies that used both the indirect and direct method of estimating LDL-C because the correlation between the two methods has been reported to be between 0.94 and 0.99 (37). However, in our analysis, the different methods may have been a source of between-study heterogeneity and this possibility should be explored further. Future investigations should also compare different doses of algal oil in the same population and compare algal oil to fish oil with cardiovascular outcomes.

In conclusion, DHA supplementation from algal oil, a marine source of (n-3) fatty acids not extracted from fish, may reduce serum TG and increase LDL-C and HDL-C in persons without CHD.




FIGURE 4 Dose of algal DHA supplementation in relation to the change in serum TG (mmol/L). Mean treatment effect and 95% CI given; n = 11.

Acknowledgments

A.M.B. and E.B.R. designed and conducted the research; A.M.B., E.L.D., W.C.W., and E.B.R. analyzed data; A.M.B. wrote the paper; and E.B.R. had primary responsibility for final content. All authors read and approved the final manuscript.

Literature Cited

- USDA. 2005 Dietary Guidelines [cited 2010 Jul 6]. Available from: <http://www.mypyramid.gov/guidelines/index.html>.
- Sacks FM, Campos H. Dietary therapy in hypertension. *N Engl J Med*. 2010;362:2102–12.
- Kris-Etherton PM, Harris WS, Appel LJ. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation*. 2002;106:2747–57.
- Lee JH, O'Keefe JH, Lavie CJ, Marchioli R, Harris WS. Omega-3 fatty acids for cardioprotection. *Mayo Clin Proc*. 2008;83:324–32.
- Lavie CJ, Milani RV, Mehra MR, Ventura HO. Omega-3 polyunsaturated fatty acids and cardiovascular diseases. *J Am Coll Cardiol*. 2009;54:585–94.
- FAO of the United Nations. The state of the world's fisheries and aquaculture; 2008 [cited 2010 Jul 6]. Available from: <http://www.fao.org/fishery/sofia/en>.
- Gussow JD. Mediterranean diets: are they environmentally responsible? *Am J Clin Nutr*. 1995;61:1383S–9S.
- Jenkins DJ, Sievenpiper JL, Pauly D, Sumaila UR, Kendall CW, Mowat FM. Are dietary recommendations for the use of fish oils sustainable? *CMAJ*. 2009;180:633–7.
- Naylor RL, Goldburg RJ, Primavera JH, Kautsky N, Beveridge MC, Clay J, Folke C, Lubchenco J, Mooney H, Troell M. Effect of aquaculture on world fish supplies. *Nature*. 2000;405:1017–24.
- Reijnders L, Soret S. Quantification of the environmental impact of different dietary protein choices. *Am J Clin Nutr*. 2003;78:664S–8S.
- Metz JG, Roessler P, Facciotti D, Levering C, Dittrich F, Lassner M, Valentine R, Lardizabal K, Domergue F, Yamada A, et al. Production of polyunsaturated fatty acids by polyketide synthases in both prokaryotes and eukaryotes. *Science*. 2001;293:290–3.
- Simopoulos AP. Omega-3 fatty acids in health and disease and in growth and development. *Am J Clin Nutr*. 1991;54:438–63.
- Gebauer SK, Psota TL, Harris WS, Kris-Etherton PM. n-3 Fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. *Am J Clin Nutr*. 2006;83:1526S–35S.
- Sijtsma L, de Swaaf ME. Biotechnological production and applications of the omega-3 polyunsaturated fatty acid docosahexaenoic acid. *Appl Microbiol Biotechnol*. 2004;64:146–53.
- Ryan AS, Keske MA, Hoffman JP, Nelson EB. Clinical overview of algal-docosahexaenoic acid: effects on triglyceride levels and other cardiovascular risk factors. *Am J Ther*. 2009;16:183–92.
- Holub BJ. Docosahexaenoic acid (DHA) and cardiovascular disease risk factors. *Prostaglandins Leukot Essent Fatty Acids*. 2009;81:199–204.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;151:264–9, W64.
- Conquer JA, Holub BJ. Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects. *J Nutr*. 1996;126:3032–9.
- Sanders TA, Gleason K, Griffin B, Miller GJ. Influence of an algal triacylglycerol containing docosahexaenoic acid (22: 6n-3) and docosapentaenoic acid (22: 5n-6) on cardiovascular risk factors in healthy men and women. *Br J Nutr*. 2006;95:525–31.
- Kelley DS, Siegel D, Vemuri M, Chung GH, Mackey BE. Docosahexaenoic acid supplementation decreases remnant-like particle-cholesterol and increases the (n-3) index in hypertriglyceridemic men. *J Nutr*. 2008;138:30–5.
- Theobald HE, Goodall AH, Sattar N, Talbot DC, Chowienzyk PJ, Sanders TA. Low-dose docosahexaenoic acid lowers diastolic blood pressure in middle-aged men and women. *J Nutr*. 2007;137:973–8.
- Stark KD, Holub BJ. Differential eicosapentaenoic acid elevations and altered cardiovascular disease risk factor responses after supplementation with docosahexaenoic acid in postmenopausal women receiving and not receiving hormone replacement therapy. *Am J Clin Nutr*. 2004;79:765–73.
- Kelley DS, Siegel D, Vemuri M, Mackey BE. Docosahexaenoic acid supplementation improves fasting and postprandial lipid profiles in hypertriglyceridemic men. *Am J Clin Nutr*. 2007;86:324–33.
- Arterburn LM, Oken HA, Hoffman JP, Bailey-Hall E, Chung G, Rom D, Hamersley J, McCarthy D. Bioequivalence of Docosahexaenoic acid from different algal oils in capsules and in a DHA-fortified food. *Lipids*. 2007;42:1011–24.
- Jensen CL, Maude M, Anderson RE, Heird WC. Effect of docosahexaenoic acid supplementation of lactating women on the fatty acid composition of breast milk lipids and maternal and infant plasma phospholipids. *Am J Clin Nutr*. 2000;71:292S–9S.
- Theobald HE, Chowienzyk PJ, Whittall R, Humphries SE, Sanders TA. LDL cholesterol-raising effect of low-dose docosahexaenoic acid in middle-aged men and women. *Am J Clin Nutr*. 2004;79:558–63.
- Vidgren HM, Agren JJ, Schwab U, Rissanen T, Hanninen O, Uusitupa MI. Incorporation of n-3 fatty acids into plasma lipid fractions, and erythrocyte membranes and platelets during dietary supplementation with fish, fish oil, and docosahexaenoic acid-rich oil among healthy young men. *Lipids*. 1997;32:697–705.
- Maki KC, Van Elsland ME, McCarthy D, Hess SP, Veith PE, Bell M, Subbiah P, Davidson MH. Lipid responses to a dietary docosahexaenoic acid supplement in men and women with below average levels of high density lipoprotein cholesterol. *J Am Coll Nutr*. 2005;24:189–99.
- Davidson MH, Maki KC, Kalkowski J, Schaefer EJ, Torri SA, Drennan KB. Effects of docosahexaenoic acid on serum lipoproteins in patients with combined hyperlipidemia: a randomized, double-blind, placebo-controlled trial. *J Am Coll Nutr*. 1997;16:236–43.
- Nelson GJ, Schmidt PC, Bartolini GL, Kelley DS, Kyle D. The effect of dietary docosahexaenoic acid on plasma lipoproteins and tissue fatty acid composition in humans. *Lipids*. 1997;32:1137–46.
- Geppert J, Kraft V, Demmelmaier H, Koletzko B. Docosahexaenoic acid supplementation in vegetarians effectively increases omega-3 index: a randomized trial. *Lipids*. 2005;40:807–14.
- Wien M, Rajaram S, Oda K, Sabate J. Decreasing the linoleic acid to alpha-linolenic acid diet ratio increases eicosapentaenoic acid in erythrocytes in adults. *Lipids*. 2010;45:683–92.
- Agren JJ, Hanninen O, Julkunen A, Fogelholm L, Vidgren H, Schwab U, Pynnönen O, Uusitupa M. Fish diet, fish oil and docosahexaenoic acid rich oil lower fasting and postprandial plasma lipid levels. *Eur J Clin Nutr*. 1996;50:765–71.
- Geppert J, Kraft V, Demmelmaier H, Koletzko B. Microalgal docosahexaenoic acid decreases plasma triacylglycerol in normolipidaemic vegetarians: a randomised trial. *Br J Nutr*. 2006;95:779–86.
- Wu WH, Lu SC, Wang TF, Jou HJ, Wang TA. Effects of docosahexaenoic acid supplementation on blood lipids, estrogen metabolism, and in vivo oxidative stress in postmenopausal vegetarian women. *Eur J Clin Nutr*. 2006;60:386–92.
- Neff LM, Culiner J, Cunningham-Rundles S, Seidman C, Meehan D, Maturi J, Wittkowski KM, Levine B, Breslow JL. Algal docosahexaenoic acid affects plasma lipoprotein particle size distribution in overweight and obese adults. *J Nutr*. 2011;141:207–13.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502.
- Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, Brown L, Warnica JW, Arnold JM, Wun CC, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med*. 1996;335:1001–9.
- Moher D, Pham B, Jones A, Cook DJ, Jadad AR, Moher M, Tugwell P, Klassen TP. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet*. 1998;352:609–13.
- Egger MS, George D, Altman DG. Systematic reviews in health care: meta-analysis in context. London: BMJ Publishing Group; 2001.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557–60.
- Eslick GD, Howe PR, Smith C, Priest R, Bensoussan A. Benefits of fish oil supplementation in hyperlipidemia: a systematic review and meta-analysis. *Int J Cardiol*. 2009;136:4–16.
- Lichtenstein AH, Appel LJ, Brands M, Carnethon M, Daniels S, Franch HA, Franklin B, Kris-Etherton P, Harris WS, Howard B, et al. Summary

- of American Heart Association Diet and Lifestyle Recommendations revision 2006. *Arterioscler Thromb Vasc Biol.* 2006;26:2186–91.
44. Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: evaluating the risks and the benefits. *JAMA.* 2006;296:1885–99.
 45. Simopoulos AP, Leaf A, Salem N Jr. Workshop on the essentiality of and recommended dietary intakes for omega-6 and omega-3 fatty acids. *J Am Coll Nutr.* 1999;18:487–9.
 46. Institute of Medicine. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids, 2002/2005 [cited 2011 Sept 4]. Available from: <http://www.iom.edu/Global/News%20Announcements/~media/C5CD2DD7840544979A549EC47E56A02B.ashx>.
 47. USDA Agricultural Research Center. National Health and Nutrition Examination Survey (NHANES) 2005–2006. What we eat in America [cited 2010 Jul 12]. Available from: http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/0506/Table_1_NIF_05.pdf.
 48. FAO of the United Nations. General situation of world fish stocks; 2002 [cited 2010 Aug 5]. Available from: <http://www.fao.org/newsroom/common/ecg/1000505/en/stocks.pdf>.
 49. Lee JH, O'Keefe JH, Lavie CJ, Harris WS. Omega-3 fatty acids: cardiovascular benefits, sources and sustainability. *Nat Rev Cardiol.* 2009;6:753–8.
 50. Martek Biosciences Corporation [cited 2010 Jul 14]. Available from: <http://www.lifesdha.com/Products-Containing-lifesDHA-/Partner-Products/tabid/683/Default.aspx>.
 51. Grimsgaard S, Bonna KH, Hansen JB, Nordoy A. Highly purified eicosapentaenoic acid and docosahexaenoic acid in humans have similar triacylglycerol-lowering effects but divergent effects on serum fatty acids. *Am J Clin Nutr.* 1997;66:649–59.
 52. Mori TA, Burke V, Puddey IB, Watts GF, O'Neal DN, Best JD, Beilin LJ. Purified eicosapentaenoic and docosahexaenoic acids have differential effects on serum lipids and lipoproteins, LDL particle size, glucose, and insulin in mildly hyperlipidemic men. *Am J Clin Nutr.* 2000;71:1085–94.
 53. Roche HM, Gibney MJ. Effect of long-chain n-3 polyunsaturated fatty acids on fasting and postprandial triacylglycerol metabolism. *Am J Clin Nutr.* 2000;71:232S–7S.
 54. Abbey M, Clifton P, Kestin M, Belling B, Nestel P. Effect of fish oil on lipoproteins, lecithin:cholesterol acyltransferase, and lipid transfer protein activity in humans. *Arteriosclerosis.* 1990;10:85–94.
 55. Berneis KK, Krauss RM. Metabolic origins and clinical significance of LDL heterogeneity. *J Lipid Res.* 2002;43:1363–79.
 56. Friedberg CE, Janssen MJ, Heine RJ, Grobbee DE. Fish oil and glycemic control in diabetes. A meta-analysis. *Diabetes Care.* 1998;21:494–500.
 57. Harris WS. n-3 fatty acids and serum lipoproteins: human studies. *Am J Clin Nutr.* 1997;65:1645S–54S.
 58. Rawn DF, Breakell K, Verigin V, Nicolidakis H, Sit D, Feeley M, Ryan JJ. Persistent organic pollutants in fish oil supplements on the Canadian market: polychlorinated dibenzo-p-dioxins, dibenzofurans, and polybrominated diphenyl ethers. *J Food Sci.* 2009;74:T31–6.
 59. Rawn DF, Breakell K, Verigin V, Nicolidakis H, Sit D, Feeley M. Persistent organic pollutants in fish oil supplements on the Canadian market: polychlorinated biphenyls and organochlorine insecticides. *J Food Sci.* 2009;74:T14–9.
 60. Amazon.com. Health and nutrition: vegetarian DHA [cited 2010 Jul 14]. Available from: http://www.amazon.com/s/qid=1279119763/ref=sr_st?keywords=vegetarian+DHA&page=1&rh=n%3A3760901%2Cn%3A13760931%2Ck%3Avegetarian+DHA&sort=price.
 61. Amazon.com. Health and personal care: diet and nutrition: fish oil EPA [cited 2010 Jul 14]. Available from: http://www.amazon.com/s/qid=1279120356/ref=sr_st?keywords=fish+oil+EPA&page=1&rh=n%3A3760901%2Cn%3A13760931%2Cn%3A3764441%2Ck%3Afish+oil+EPA&sort=price.